NUTRITION, FOOD QUALITY AND EXPOSURE REDUCTION IN ALS

A practical, evidence-informed paper on eating well, limiting heavily processed foods, and protecting nutritional status while living with amyotrophic lateral sclerosis

Purpose
To explain how thoughtful food choices can support nutrition, hydration, weight maintenance and day-to-day well-being in ALS, while offering a realistic approach to reducing avoidable dietary exposures without creating unnecessary fear or restrictive eating.

Educational resource • United States • November 2026

This paper is for education and discussion with an ALS care team. It is not a personalized medical nutrition prescription.

Important perspective: nutrition in ALS is not an ordinary weight-loss project

ALS can make eating more difficult at the same time that the body may have substantial energy needs. Chewing can become tiring, swallowing may become less safe, meals can take longer, and weakness in the arms or hands can make shopping, cooking and feeding oneself harder. Some people also experience changes in appetite, constipation, fatigue or breathing that complicate meals.

For these reasons, the first nutritional priority in ALS is usually to maintain adequate intake, hydration and an appropriate body weight—not to pursue a restrictive diet, cleanse, detox program, or idealized version of “clean eating.” Unintentional weight loss and inadequate intake can worsen nutritional status and contribute to loss of muscle and energy. The ALS Association emphasizes sufficient calories, protein, fluids and individualized texture adjustments. [1,2]

The central rule: Improve food quality where it is practical, but do not allow food rules to reduce calories, protein, fluids, enjoyment or safe swallowing. If a “healthier” substitution causes weight loss or makes eating harder, it may not be the healthier choice for that person.

There is no specific food, supplement, eating pattern or list of forbidden ingredients that has been shown to cure ALS or reliably stop its progression. The ALS Association’s swallowing and nutrition guide states that no particular foods or beverages have been shown to improve ALS outcomes and none should be universally avoided. Individual needs differ. [3] This paper therefore distinguishes well-established general nutrition principles from hypotheses and areas where evidence is incomplete.

1. Why eating properly matters

Adequate energy and weight maintenance

Calories are energy. The body uses them to maintain temperature, support organs, power movement and breathing, repair tissues and carry out everyday functions. In ALS, energy requirements vary from person to person and can change over time. Some people may expend more energy than expected; others may have lower needs as activity and body composition change. A dietitian can help estimate needs and adjust them using weight trends, intake, symptoms and, when available, metabolic assessment.

Weight is a useful practical signal. A downward trend may mean that food intake is not keeping up with needs, even if meals appear substantial. Weighing under consistent conditions and reviewing the trend with the care team is more informative than reacting to one measurement. Do not begin a weight-loss diet or intentionally restrict calories without discussing it with the ALS clinician and dietitian.

Protein and muscle maintenance

Protein supplies amino acids used to build and maintain muscle and other body tissues. ALS itself affects motor neurons and muscle function; food cannot reverse that neurological process. Still, inadequate protein and calories can add preventable nutritional stress. Include protein sources regularly when safe and acceptable: eggs, fish, poultry, meat, milk, yogurt, cottage cheese, cheese, tofu, beans, lentils, nut or seed butters, and nutrition drinks when recommended.

When appetite is low or chewing is tiring, protein can be incorporated into foods already being eaten: yogurt blended into smoothies, powdered milk mixed into suitable foods, soft scrambled eggs, tender fish with sauce, or pureed beans. Texture and thickness must match the individual's swallowing plan.

Micronutrients, fiber and overall dietary variety

Vegetables, fruits, whole grains, legumes, nuts, seeds and dairy or fortified alternatives provide vitamins, minerals, fiber and other food components. A varied diet helps cover nutritional needs and avoids relying on a narrow range of foods. Fiber may help bowel regularity, but high-fiber foods can be difficult to chew or swallow for some people. Cook, mash, peel, blend or choose softer alternatives as needed.

Hydration

Fluids support circulation, digestion, temperature regulation and bowel function. Dehydration can contribute to constipation, dizziness, confusion and fatigue. In ALS, swallowing thin liquids may become difficult; simply drinking more without a swallowing assessment may be unsafe for someone with dysphagia. Ask a speech-language pathologist (SLP) and dietitian what fluid thickness, delivery method and daily target are appropriate.

Immune and general health

Adequate energy, protein and micronutrients support normal immune function and recovery from everyday illness. Food safety also matters: a foodborne infection can be especially disruptive when a person already has limited respiratory reserve, fatigue or difficulty clearing secretions. Good hand hygiene, careful storage, thorough cooking and avoiding cross-contamination are sensible precautions.

2. What “processed food” means—and what deserves attention

Processing is not a simple good-versus-bad category. Freezing vegetables, canning beans, pasteurizing milk, milling oats, fermenting yogurt, freezing fish and cooking food are all forms of processing. These can make nutritious foods safer, more affordable, easier to store or easier to prepare.

The concern is more often a pattern dominated by highly or ultra-processed products that are convenient but frequently contain combinations of refined starches, added sugars, excess sodium, low-fiber ingredients, saturated fats, flavorings and preservatives. Examples can include sugary drinks, candy, packaged snack cakes, many chips, some instant meals, processed meat products and some ready-to-eat meals. The category is broad, and not every packaged product has the same nutritional profile.

Why a diet dominated by highly processed products can be a problem

- **Low nutrient density:** Some products provide many calories but relatively little protein, fiber or micronutrient variety.
- **Added sugars:** Sweetened beverages, desserts and sweet snacks can displace foods that provide protein, fiber and micronutrients.
- **High sodium:** Frequent high-sodium meals and snacks can make it harder to follow individualized blood-pressure or fluid-management advice.
- **Refined carbohydrates:** Products made largely from refined flour or starch may be less filling and contain less fiber than intact or minimally processed grain choices.
- **Processed meats:** Bacon, ham, hot dogs, sausages and deli meats often contain substantial sodium; the World Health Organization's cancer agency concludes that processed meat consumption causes colorectal cancer. This is a long-term population-level cancer finding, not evidence that an occasional serving worsens ALS. [4]
- **Ease of over-reliance:** When cooking is difficult, packaged convenience foods can become the default. The goal is not to ban them, but to build a workable mix of convenient foods that also deliver nutrition.

A person with ALS may need calorie-dense foods, including foods that conventional weight-management advice would label high in fat or calories. Olive oil, avocado, nut butter, full-fat yogurt, cheese, eggs or enriched shakes may be useful

when weight maintenance is a priority. Avoid applying generic low-fat or low-calorie rules without individualized advice.

3. Common food additives and chemical exposures

The word “chemical” can sound alarming, but all food—including water, fruit and vegetables—is made of chemicals. A chemical name on a label does not by itself establish danger. Risk depends on the substance, dose, route and duration of exposure, the person's circumstances and the quality of the evidence. The FDA explains that a meaningful safety assessment considers how much of a substance is present and how much people actually consume. [5]

The aim should be sensible exposure reduction—not trying to eliminate every chemical or treating every additive as toxic. The following are examples people may encounter in food or food production.

Sodium nitrite and nitrate

These compounds are used in some cured or processed meats to preserve color, inhibit certain bacteria and contribute to flavor. Under some conditions, nitrites can participate in the formation of nitrosamines, a group of compounds that includes substances of health concern. Processed meat itself is associated with colorectal cancer risk. A practical step is to make bacon, hot dogs, salami, bologna and similar cured meats occasional rather than everyday protein choices, replacing them when feasible with eggs, fish, poultry, beans or other tolerated proteins.

Added preservatives and antioxidants

Ingredients such as sodium benzoate, potassium sorbate, sulfites, BHA and BHT may be used to inhibit spoilage or oxidation in particular products. Their permitted uses and safety assessments differ. Some individuals have specific sensitivities—for example, sulfites can trigger reactions in susceptible people—but the presence of a preservative does not mean a product is unsafe for everyone. If a clinician has identified a specific allergy or sensitivity, follow that advice. Otherwise, prioritize the overall food pattern rather than attempting to eliminate every preservative.

Artificial colors, flavors and sweeteners

Color additives, flavorings and non-sugar sweeteners are used for appearance, taste or formulation. They are not nutritionally necessary in most foods, but their presence alone does not establish that a food is harmful. If a product relies heavily on sweeteners or flavorings and provides little nutritional value, choosing a more nutrient-dense alternative may be reasonable. Do not assume that “natural” flavoring is automatically safer or that an unfamiliar ingredient is automatically dangerous.

Emulsifiers, stabilizers and texture agents

Ingredients such as lecithins, gums, carrageenan and certain modified starches help packaged foods remain mixed, creamy or stable. Research into possible effects of some additives on the gut microbiome and inflammation is evolving, much of it not sufficient to prove that ordinary food-level exposure causes disease in humans or affects ALS. For now, it is reasonable to reduce dependence on highly formulated snack foods when easy alternatives exist, but there is no ALS-specific evidence supporting a blanket ban on these ingredients.

Pesticide residues

Pesticides are used to protect crops. Residues may remain on some foods, and exposure can occur through food, water and other routes. Washing produce under clean running water and scrubbing firm surfaces can reduce dirt, microbes and some surface residues, but washing does not remove all pesticide residues. Peeling or trimming can reduce some residues but may also remove fiber and nutrients. Eating a variety of fruits and vegetables from different sources is a practical way to avoid relying on a narrow range of foods. Organic produce may be a personal choice, but organic farming can also use approved pesticides and organic food still needs washing. [6]

Food-contact substances: BPA, phthalates and PFAS

Some chemicals are associated with packaging or materials that contact food. Bisphenols such as BPA have been used in certain plastics and can linings; phthalates have had uses as plasticizers in some food-contact applications; and PFAS have been used in certain grease-resistant materials and other products. Uses and regulatory status change over time, and exposure varies by substance and source. FDA information describes food-contact substances and its ongoing evaluation of specific chemicals. The FDA reports that most foods in the general U.S. food supply tested to date have not had detectable PFAS, while noting that foods from specific contaminated areas may require attention. [7,8]

Low-effort precautions include using glass, ceramic or stainless-steel containers for hot foods when convenient; avoiding heating food in containers not labeled microwave-safe; not reusing damaged plastic containers; and following package directions. These steps are precautionary exposure-reduction habits, not proof that ordinary packaged food is unsafe.

Acrylamide and other cooking-related contaminants

Acrylamide can form naturally when some starchy plant foods—such as potatoes and grain products—are cooked at high temperatures, especially frying, roasting or baking. It is more likely to accumulate with higher temperatures and longer cooking. FDA notes that high-dose animal studies raise concern, while the human risk at typical food exposure remains uncertain. If desired, reduce frequent intake of heavily browned fries, chips and toast; favor boiling or steaming more often; and cook potatoes and toast to golden rather than dark brown. Do not avoid nutritious whole grains because of acrylamide concerns. [9]

Other process contaminants can form during certain high-temperature cooking or processing methods. Charring meat over an open flame can produce compounds such as heterocyclic amines and polycyclic aromatic hydrocarbons. Avoid routinely burning or blackening foods; use gentler cooking methods such as steaming, simmering, baking at moderate temperatures or stewing.

Sodium, added sugars and industrial trans fats

These are not all “chemicals” in the same sense, but they are common features of less healthful dietary patterns. Read the Nutrition Facts label for sodium and added sugars. Partially hydrogenated oils were the major source of artificial trans fat in the U.S. food supply and are no longer generally permitted in foods, although checking labels and imported products remains sensible. Do not confuse naturally occurring fats with industrial trans fats. For ALS, calorie needs may outweigh a generic goal of minimizing fat; focus on the type of food, the person's needs and clinician guidance.

4. Building a nourishing, practical food pattern

A helpful foundation is a flexible mix of foods that are nutrient-rich, familiar, affordable and safe to swallow. A “whole-food” approach means emphasizing foods close to their original form when practical—not refusing all canned, frozen, fortified or prepared foods.

Protein choices

- Eggs, soft omelets or scrambled eggs; fish; tender poultry or meat with sauce; yogurt, cottage cheese and milk; tofu; lentils and beans; nut or seed butters when texture-safe.
- Use moist preparations—soups, stews, casseroles, sauces and gravies—to reduce dryness and chewing effort.
- If protein intake is low, ask a dietitian whether a high-protein oral nutrition supplement or food fortification is appropriate.

Vegetables and fruits

- Choose fresh, frozen or canned options according to cost, convenience, tolerance and swallowing safety. Frozen vegetables without heavy sauces and canned beans or fruit can be practical choices.
- Cook vegetables until soft; mash or puree if needed. Soft fruit, applesauce, ripe banana and canned fruit may be easier than raw, fibrous produce.
- Add vegetables or fruit to soups, smoothies, oatmeal, yogurt or sauces to increase variety without making meals difficult.

Whole grains, legumes and starches

- Oatmeal, soft whole-grain cereals, rice, pasta, potatoes, quinoa, soft breads and beans can provide energy and fiber.
- Choose the texture that is safe. A dry crusty whole-grain product is not automatically a better choice if it is hard to chew or swallow.

Fats and calorie enrichment

Olive oil, avocado, nut butters, seeds, dairy fats, mayonnaise and dressings can increase calories in a small volume. For someone losing weight or tiring before finishing meals, adding calories to foods already tolerated may be more useful than increasing the amount of food on the plate. A dietitian can help select additions that fit medical conditions, preferences and swallowing needs.

Convenience foods that can still be good choices

Convenience is not the enemy. Consider frozen vegetables, canned beans rinsed if sodium is a concern, plain frozen fish, pre-cooked eggs, plain yogurt, unsweetened applesauce, oatmeal, low-sodium soups, rotisserie chicken if safely prepared and tolerated, and oral nutrition drinks recommended by the care team. If a packaged meal is the only realistic option on a difficult day, eating enough is more important than achieving a perfect ingredient list.

5. Reducing avoidable exposures: a realistic kitchen plan

- **Read labels selectively.** Compare serving size, protein, fiber, sodium and added sugars. Scan the ingredient list for the main ingredients rather than trying to interpret every unfamiliar word.
- **Choose variety.** Rotate fruits, vegetables, grains and protein sources. Variety supports nutrient coverage and reduces dependence on one food or source.
- **Wash produce under running water.** Rub produce with your hands; scrub firm produce with a clean brush. Do not use dish soap, detergent or household cleaners on food. FDA and USDA guidance advises against washing produce with soap. [10]
- **Handle food safely.** Wash hands and surfaces; separate raw meat, poultry and seafood from ready-to-eat foods; refrigerate perishable foods promptly; cook foods to safe temperatures; and discard food with signs of spoilage.
- **Use gentler cooking methods often.** Steam, poach, simmer, stew or bake without charring. Avoid routinely deep-frying or blackening foods. Do not let concern about cooking contaminants prevent adequate food intake.
- **Use sensible food storage.** Store leftovers promptly in shallow containers and refrigerate. Reheat thoroughly. When in doubt about time or temperature, discard questionable food rather than taking a risk.
- **Limit routine processed-meat intake.** Use fresh or minimally processed protein options more often when practical, but do not skip meals if a familiar processed food is the only food that is manageable.

- **Make packaging changes easy.** Use glass or ceramic for reheating when practical, and do not microwave food in packaging unless the manufacturer says it is microwave-safe.
- **Do not spend beyond your means.** Organic, specialty or premium foods are not required for good nutrition. Frozen, canned and conventional produce can all contribute to a nourishing diet.

6. ALS-specific considerations: safety, effort and swallowing

Swallowing changes require professional assessment

Coughing or choking during meals, a wet or gurgly voice after swallowing, food sticking in the throat, unexplained weight loss, meals taking much longer, recurrent chest infections or difficulty managing saliva should be reported promptly to the ALS team. Dysphagia can lead to aspiration—food or liquid entering the airway—and aspiration can cause serious lung complications. A speech-language pathologist can assess swallowing and recommend appropriate textures, liquid thickness, posture, pacing and strategies. Do not independently adopt a thickened-liquid or chin-tuck strategy without individualized advice.

Texture modification should preserve nutrition

Soft, minced, moist or pureed foods may be safer or easier for some people, but texture changes can unintentionally reduce calories and variety. A dietitian can show how to enrich purees and soft meals with oils, dairy, protein powders or other ingredients that fit the swallowing plan. Foods that are dry, crumbly, sticky, stringy or mixed in texture may be difficult for some people; the individual's SLP should identify specific risks.

Fatigue and mealtime effort

- Try smaller, more frequent meals or snacks rather than relying on three large meals.
- Eat when energy is best; consider resting before meals and allowing enough time.
- Use adaptive utensils, supportive seating and caregiver assistance when helpful.
- Prepare easy-to-eat foods in advance or ask for help with shopping, cutting and meal preparation.
- Avoid talking or multitasking during meals if the swallowing team recommends focused, unhurried eating.

Feeding tubes and nutrition support

A feeding tube is a medical decision, not a failure and not a reason to stop enjoying food by mouth if oral intake remains safe. It can help deliver calories, fluids and medication when eating becomes too difficult or unsafe. Discuss the potential benefits, burdens, timing and procedural considerations early with the ALS team, rather than waiting for a crisis. The ALS Association and clinical nutrition guidelines recommend ongoing monitoring and individualized discussion. [2,3,11]

Supplements and “detox” products

A supplement is not automatically safe because it is natural, and high-dose vitamins or herbal products can interact with medications or cause harm. There is insufficient evidence that routine high-dose vitamin, antioxidant or “detox” regimens improve ALS outcomes. Use supplements to address a documented or likely nutritional gap under professional guidance. Bring the full list of supplements, powders, oils and nonprescription products to appointments. Avoid products promising to cleanse the body, remove toxins or treat ALS.

7. A practical daily approach

The following is a planning framework, not a fixed menu or calorie prescription. Each item must be adapted to the person's swallowing assessment, allergies, preferences, medical conditions and energy needs.

Eating occasion	Flexible example	Nutrition purpose
Breakfast	Oatmeal enriched with milk; soft eggs; yogurt or a smoothie if safe	Energy, protein and a manageable texture
Mid-morning	Yogurt, pudding, applesauce with nut butter if texture-safe, or a prescribed nutrition drink	Additional calories and protein
Lunch	Moist chicken, fish, tofu or beans with soft vegetables and rice or potato	Protein, starch and dietary variety
Afternoon	Smoothie, cheese and soft fruit, or another tolerated snack	Prevent long gaps and reduce meal burden
Dinner	Stew, casserole, tender protein, cooked vegetables and a preferred starch	Nourishment in a moist, adaptable form
Evening	Milk, yogurt, fortified pudding or snack if needed	Help meet daily needs

The right daily plan may include more calories, more fat, more snacks or more convenience foods than a standard public-health menu. That is not a contradiction. The person’s actual nutritional needs and safe swallowing plan take priority.

8. What to monitor and discuss with the ALS team

- Weight trend and whether clothing is becoming looser; report unplanned weight loss early.
- How long meals take, whether eating causes exhaustion, and whether appetite or food enjoyment has changed.
- Coughing, choking, wet voice, food sticking, drooling, recurrent respiratory infections or changes in swallowing.
- Daily fluid intake, urine changes, constipation and signs of dehydration.
- Whether hand weakness, posture, seating, utensils or caregiver availability are limiting food intake.
- Medication and supplement list, including possible interactions and whether medications can be taken safely with modified textures.
- Whether a dietitian, SLP, occupational therapist or feeding-tube consultation would help.

Many ALS resources recommend regular nutritional monitoring. The ALS Association guide discusses weight maintenance and individualized swallowing support; a clinical guideline recommends monitoring weight and nutritional status at regular follow-up intervals. The schedule should be set by the treating team and adjusted when symptoms change. [3,11]

9. Common mistakes to avoid

- **Over-restricting.** Eliminating many foods to avoid every additive, pesticide residue or processed ingredient can shrink the diet and contribute to inadequate intake.

- **Confusing “natural” with safe.** Natural products can contain allergens, contaminants or pharmacologically active ingredients.
- **Assuming organic means chemical-free.** Organic production can use approved pesticides; washing is still important.
- **Using weight-loss advice designed for the general population.** Calorie restriction may be inappropriate in ALS, especially with unplanned weight loss.
- **Ignoring swallowing changes.** A nutritious food is not beneficial if it is unsafe to swallow.
- **Starting supplements without review.** High doses and multi-ingredient products can create interactions or toxicity.
- **Believing one food causes or cures ALS.** ALS is complex. Food choices can support general health and nutritional status, but claims that a specific ingredient causes ALS or that a diet reverses it go beyond established evidence.

10. Conclusion: prioritize nourishment, then improve quality

A thoughtful diet for someone living with ALS should be nourishing, flexible, safe to swallow, practical to prepare and enjoyable. Emphasizing a variety of minimally processed foods can improve the overall nutritional quality of the diet and reduce reliance on products high in added sugars, sodium and processed meats. Washing produce, using sensible food-storage practices, avoiding frequent charring and making practical packaging choices can reduce some avoidable exposures.

But the most important principle is balance. Reducing exposure to unwanted ingredients should never become a reason to eat too little, lose weight unintentionally, become dehydrated or give up foods that make life more enjoyable. The healthiest plan is the one that meets the person's needs and can be sustained.

For ALS, the practical priorities are: maintain an appropriate weight; consume adequate calories and protein; stay hydrated using swallowing-safe liquids; monitor changes early; choose varied foods; reduce heavily processed foods where practical; and work with an ALS dietitian and speech-language pathologist as needs evolve.

References and further reading

- 1. The ALS Association. “Nutritional Guidance for ALS.” Practical information on protein, fiber, hydration, calories and swallowing-related adaptations. https://www.als.org/navigating-als/living-with-als/therapies-care/nutrition-feeding-tubes
- 2. The ALS Association. “Maintaining Adequate Nutrition: A Continuing Challenge in ALS.” Discussion of energy needs, weight maintenance, hydration and nutrition support. https://www.als.org/navigating-als/living-als/therapies-and-care/nutrition-and-feeding-tubes/maintaining-adequate
- 3. The ALS Association. Guide #8, “Adjusting to Swallowing Changes and Nutritional Management in ALS” (2025 resource guide). https://www.als.org/sites/default/files/2025-04/Resource-Guide-8.pdf
- 4. World Health Organization / International Agency for Research on Cancer. “Carcinogenicity of the consumption of red meat and processed meat.” https://www.who.int/news-room/questions-and-answers/item/cancer-carcinogenicity-of-the-consumption-of-red-meat-and-processed-meat/
- 5. U.S. Food and Drug Administration. “Is Food Safe if it Has Chemicals?” Explanation of dose, exposure and risk. https://www.fda.gov/consumers/consumer-updates/food-safe-if-it-has-chemicals
- 6. U.S. Environmental Protection Agency. “Pesticides and Food: Healthy, Sensible Food Practices.” https://www.epa.gov/safepestcontrol/pesticides-and-food-healthy-sensible-food-practices
- 7. U.S. Food and Drug Administration. “Food Packaging & Other Substances that Come in Contact with Food.” https://www.fda.gov/food/food-ingredients-packaging/food-packaging-other-substances-come-contact-food-information-consumers

- 8. U.S. Food and Drug Administration. “Questions and Answers on PFAS in Food.” https://www.fda.gov/food/process-contaminants-food/questions-and-answers-pfas-food
- 9. U.S. Food and Drug Administration. “Acrylamide and Diet, Food Storage, and Food Preparation.” https://www.fda.gov/food/process-contaminants-food/acrylamide-and-diet-food-storage-and-food-preparation
- 10. U.S. Food and Drug Administration. “Selecting and Serving Produce Safely.” Guidance on washing and handling fruits and vegetables. https://www.fda.gov/food/buy-store-serve-safe-food/selecting-and-serving-produce-safely
- 11. Burgos, R. et al. ESPEN guideline clinical nutrition in neurology. Clinical Nutrition. 2018;37:354–396. Includes recommendations for dysphagia screening, nutritional monitoring and adapting/enriching meals. https://www.espen.org/files/ESPEN-Guidelines/ESPEN-guideline_clinical_nutrition_in_neurology.pdf
- 12. Evidence-Based Nutritional Recommendations for Maintaining or Restoring Nutritional Status in Patients with ALS: A Systematic Review. 2025. https://pmc.ncbi.nlm.nih.gov/articles/PMC11901627/

Medical and scientific limitations

This paper summarizes selected public guidance and research available as of November 2026. It is not a systematic review and does not evaluate every food additive, contaminant, dietary pattern or ALS study. Evidence about general diet quality should not be mistaken for evidence of an ALS treatment. Individual recommendations may differ with swallowing function, weight trajectory, diabetes, kidney or heart disease, allergies, medications, feeding-tube status and personal goals. Discuss meaningful dietary changes with the ALS care team.

Prepared as an educational printable resource. Readers should consult the linked original sources for full context and updates.